

2.3 Factor Theorem Lesson with solutions

This document was composed and formatted by Annie Bernatchez.

The course material originated from www.jensenmath.ca

In this section, you will learn how to determine the factors of a polynomial function of degree 3 or greater.

Part 1: Remainder Theorem Refresher

a) Use the remainder theorem to determine the remainder when $f(x) = x^3 + 4x^2 + x - 6$ is divided by $x + 2$

$$\begin{aligned} f(-2) &= (-2)^3 + 4(-2)^2 + (-2) - 6 \\ &= -8 + 16 - 2 - 6 \\ &= 0 \end{aligned}$$

The remainder when divided by $x + 2$ is 0. This means that $x + 2$ is a factor of the dividend.

Remainder Theorem: When a polynomial function $P(x)$ is divided by $x - b$, the remainder is $P(b)$; and when it is divided by $ax - b$, the remainder is $P\left(\frac{b}{a}\right)$, where a and b are integers, and $a \neq 0$.

b) Verify your answer to part a) by completing the division using long division or synthetic division.

Note: I chose synthetic since it is a linear divisor of the form $x - b$.

$$\begin{array}{r|rrrr} -2 & 1 & 4 & 1 & -6 \\ & & -2 & -4 & 6 \\ \hline & 1 & 2 & -3 & 0 \\ & x^2 & x & \# & R \end{array}$$

$$x^3 + 4x^2 + x - 6 = (x + 2)(x^2 + 2x - 3)$$

Factor Theorem: $x - b$ is a factor of a polynomial $P(x)$ if and only if $P(b) = 0$. Similarly, $ax - b$ is a factor of $P(x)$ if and only if $P\left(\frac{b}{a}\right) = 0$.

Example 1: Determine if $x - 3$ and $x + 2$ are factors of $P(x) = x^3 - x^2 - 14x + 24$

$$\begin{aligned} P(3) &= (3)^3 - (3)^2 - 14(3) + 24 \\ &= 27 - 9 - 42 + 24 \\ &= 0 \end{aligned}$$

Since the remainder is 0, $x - 3$ divides evenly into $P(x)$; that means $x - 3$ is a factor of $P(x)$.

$$\begin{aligned} P(-2) &= (-2)^3 - (-2)^2 - 14(-2) + 24 \\ &= -8 - 4 + 28 + 24 \\ &= 40 \end{aligned}$$

Since the remainder is not 0, $x + 2$ does not divide evenly into $P(x)$; that means $x + 2$ is not a factor of $P(x)$.

Part 2: How to Determine a Factor of a Polynomial With Leading Coefficient 1

You could guess and check values of b that make $P(b) = 0$ until you find one that works...

Or you can use the Integral Zero Theorem to help.

Integral Zero Theorem: If $x - b$ is a factor of a polynomial function $P(x)$ with leading coefficient 1 and remaining coefficients that are integers, then b is a factor of the constant term of $P(x)$.

Note: Once one of the factors of a polynomial is found, division is used to determine the other factors.

Example 2: Factor $x^3 + 2x^2 - 5x - 6$ fully.

$$\text{Let } P(x) = x^3 + 2x^2 - 5x - 6$$

Find a value of b such that $P(b) = 0$. Based on the factor theorem, if $P(b) = 0$, then we know that $x - b$ is a factor. We can then divide $P(x)$ by that factor.

The integral zero theorem tells us to test factors of -6 .

2.3 Factor Theorem Lesson with solutions

Test ± 1 , ± 2 , ± 3 , and ± 6 . Once one factor is found, you can stop testing and use that factor to divide $P(x)$.

$$\begin{aligned} P(1) &= (1)^3 + 2(1)^2 - 5(1) - 6 \\ &= 1 + 2 - 5 - 6 \\ &= -8 \end{aligned}$$

Since $P(1) \neq 0$, we know that $x - 1$ is **NOT** a factor of $P(x)$.

$$\begin{aligned} P(2) &= (2)^3 + 2(2)^2 - 5(2) - 6 \\ &= 8 + 8 - 10 - 6 \\ &= 0 \end{aligned}$$

Since $P(2) = 0$, we know that $x - 2$ is a factor of $P(x)$.

You can now use either long division or synthetic division to find the other factors.

Method 1: Long division

$$\begin{array}{r} x^2 + 4x + 3 \\ x - 2 \overline{) x^3 + 2x^2 - 5x - 6} \\ \underline{x^3 - 2x^2} \\ 4x^2 - 5x \\ \underline{4x^2 - 8x} \\ 3x - 6 \\ \underline{3x - 6} \\ R = 0 \end{array}$$

Method 2: Synthetic Division

$$\begin{array}{r|rrrr} 2 & 1 & 2 & -5 & -6 \\ & & 2 & 8 & 6 \\ \hline & 1 & 4 & 3 & 0 \\ & x^2 & x & \# & R \end{array}$$

$$x^3 + 2x^2 - 5x - 6 = (x - 2)(x^2 + 4x + 3)$$

$$= (x - 2)(x + 3)(x + 1)$$

Example 3: Factor $x^4 + 3x^3 - 7x^2 - 27x - 18$ completely.

$$\text{Let } P(x) = x^4 + 3x^3 - 7x^2 - 27x - 18$$

Find a value of b such that $P(b) = 0$. Based on the factor theorem, if $P(b) = 0$, then we know that $x - b$ is a factor. We can then divide $P(x)$ by that factor.

The integral zero theorem tells us to test factors of -18 .

Test ± 1 , ± 2 , ± 3 , ± 6 , ± 9 , and ± 18 . Once one factor is found, you can stop testing and use that factor to divide $P(x)$.

$$\begin{aligned} P(1) &= (1)^4 + 3(1)^3 - 7(1)^2 - 27(1) - 18 \\ &= -48 \end{aligned}$$

$x - 1$ is **NOT** a factor of $P(x)$.

$$\begin{aligned} P(-1) &= (-1)^4 + 3(-1)^3 - 7(-1)^2 - 27(-1) - 18 \\ &= 0 \end{aligned}$$

$x + 1$ **IS** a factor of $P(x)$.

Since $P(-1) = 0$, this tells us that $x + 1$ is a factor. Use division to determine the other factor.

$$\begin{array}{r|rrrrrr} -1 & 1 & 3 & -7 & -27 & -18 \\ & & -1 & -2 & 9 & 18 \\ \hline & 1 & 2 & -9 & -18 & 0 \\ & x^3 & x^2 & x & \# & R \end{array}$$

$$x^4 + 3x^3 - 7x^2 - 27x - 18 = (x + 1)(x^3 + 2x^2 - 9x - 18)$$

We can now further divide $x^3 + 2x^2 - 9x - 18$ using division again or by factoring by grouping.

Method 1: Division

Test factors of -18

$$\begin{aligned} f(-2) &= (-2)^3 + 2(-2)^2 - 9(-2) - 18 \\ &= 0 \end{aligned}$$

$x + 2$ is a factor

$$\begin{array}{r|rrrr} -2 & 1 & 2 & -9 & -18 \\ & & -2 & 0 & 18 \\ \hline & 1 & 0 & -9 & 0 \\ & x^2 & x & \# & R \end{array}$$

$$x^4 + 3x^3 - 7x^2 - 27x - 18 = (x + 1)(x + 2)(x^2 - 9)$$

$$= (x + 1)(x + 2)(x - 3)(x + 3)$$

Method 2: Factoring by Grouping

$$f(x) = x^3 + 2x^2 - 9x - 18$$

Group the first 2 terms and the last 2 terms and separate with an addition sign.

$$f(x) = (x^3 + 2x^2) + (-9x - 18)$$

Common factor within each group

$$f(x) = x^2(x + 2) - 9(x + 2)$$

Factor out the common binomial

$$f(x) = (x + 2)(x^2 - 9)$$

Therefore,

$$x^4 + 3x^3 - 7x^2 - 27x - 18 = (x + 1)(x^3 + 2x^2 - 9x - 18)$$

$$= (x + 1)(x + 2)(x^2 - 9)$$

$$= (x + 1)(x + 2)(x - 3)(x + 3)$$

Example 4: Try Factoring by Grouping Again

$$x^4 - 6x^3 + 2x^2 - 12x$$

Note: Factoring by grouping does not always work...but when it does, it saves you time!

$$= (x^4 - 6x^3) + (2x^2 - 12x)$$

$$= x^3(x - 6) + 2x(x - 6)$$

$$= (x - 6)(x^3 + 2x)$$

$$= (x - 6)(x)(x^2 + 2)$$

Part 3: How to Determine a Factor of a Polynomial With Leading Coefficient NOT 1

The integral zero theorem can be extended to include polynomials with leading coefficients that are not 1. This extension is known as the rational zero theorem.

Rational Zero Theorem: Suppose $P(x)$ is a polynomial function with integer coefficients and $x = \frac{b}{a}$ is a zero of $P(x)$, where a and b are integers and $a \neq 0$. Then,

- b is a factor of the constant term of $P(x)$
- a is a factor of the leading coefficient of $P(x)$
- $(ax - b)$ is a factor of $P(x)$

Example 5: Factor $P(x) = 3x^3 + 2x^2 - 7x + 2$

We must start by finding a value of $\frac{b}{a}$ where $P\left(\frac{b}{a}\right) = 0$.

b must be a factor of the constant term. Possible values for b are: $\pm 1, \pm 2$

a must be a factor of the leading coefficient. Possible values of a are: $\pm 1, \pm 3$

Therefore, possible values for $\frac{b}{a}$ are: $\pm 1, \pm \frac{1}{3}, \pm 2, \pm \frac{2}{3}$

Test values of $\frac{b}{a}$ for x in $P(x)$ to find a zero.

$$P(1) = 3(1)^3 + 2(1)^2 - 7(1) + 2 = 0$$

Since $P(1) = 0$, $x - 1$ is a factor of $P(x)$. Use division to find the other factors.

$$\begin{array}{r|rrrr} 1 & 3 & 2 & -7 & 2 \\ & & 3 & 5 & -2 \\ \hline & 3 & 5 & -2 & 0 \\ & x^2 & x & \# & R \end{array}$$

$$3x^3 + 2x^2 - 7x + 2 = (x - 1)(3x^2 + 5x - 2)$$

$$= (x - 1)[(3x^2 + 6x) + (-1x - 2)]$$

$$= (x - 1)[3x(x + 2) - 1(x + 2)]$$

$$= (x - 1)(x + 2)(3x - 1)$$

Example 6: Factor $P(x) = 2x^3 + x^2 - 7x - 6$

Possible values for b are: $\pm 1, \pm 2, \pm 3, \pm 6$

Possible values of a are: $\pm 1, \pm 2$

Therefore, possible values for $\frac{b}{a}$ are: $\pm 1, \pm \frac{1}{2}, \pm 2, \pm 3, \pm \frac{3}{2}, \pm 6$

$$f(-1) = 2(-1)^3 + (-1)^2 - 7(-1) - 6 = 0$$

Therefore, $x + 1$ is a factor of $P(x)$

$$\begin{array}{r|rrrr} -1 & 2 & 1 & -7 & -6 \\ & & -2 & 1 & 6 \\ \hline & 2 & -1 & -6 & 0 \\ & x^2 & x & \# & R \end{array}$$

$$2x^3 + x^2 - 7x - 6 = (x + 1)(2x^2 - x - 6)$$

$$= (x + 1)[(2x^2 - 4x) + (3x - 6)]$$

$$= (x + 1)[2x(x - 2) + 3(x - 2)]$$

$$= (x + 1)(x - 2)(2x + 3)$$

Part 4: Application Question

Example 7: When $f(x) = 2x^3 - mx^2 + nx - 2$ is divided by $x + 1$, the remainder is -12 and $x - 2$ is a factor. Determine the values of m and n .

Hint: Use the information given to create 2 equations and then use substitution or elimination to solve.

$$\begin{aligned} f(-1) &= 2(-1)^3 - m(-1)^2 + n(-1) - 2 \\ -12 &= -2 - m - n - 2 \\ -8 &= -m - n \end{aligned}$$

Equation (1): $-8 = -m - n$

$$\begin{aligned} f(2) &= 2(2)^3 - m(2)^2 + n(2) - 2 \\ 0 &= 16 - 4m + 2n - 2 \\ -14 &= -4m + 2n \end{aligned}$$

Equation (2): $-14 = -4m + 2n$

Multiply equation (1) by 2: $-16 = -2m - 2n$

Add this to equation (2): $-14 = -4m + 2n$

$$\begin{aligned} -30 &= -6m \\ 5 &= m \end{aligned}$$

Substitute $m = 5$ into equation (1) or (2):

$$\begin{aligned} -8 &= -5 - n \\ n &= 3 \end{aligned}$$

So $m = 5$ and $n = 3$