

2.1 Long Division of Polynomials and the Remainder Theorem Worksheet

This document was composed and formatted by Annie Bernatchez.

The course material originated from www.jensenmath.ca

1) Use the remainder theorem to determine the remainder when $2x^3 + 7x^2 - 8x + 3$ is divided by each binomial.

a) $x + 1$

b) $x - 2$

c) $x + 3$

2) a) Divide $x^3 + 3x^2 - 2x + 5$ by $x + 1$. Express the result in quotient form.

b) Write the corresponding statement that can be used to check the division.

3) Divide $3x^4 - 4x^3 - 6x^2 + 17x - 8$ by $3x - 4$. Express the result in quotient form.

b) Write the corresponding statement that can be used to check the division.

4) Perform each division. Express the result in quotient form.

a) $x^3 + 7x^2 - 3x + 4$ divided by $x + 2$

b) $6x^3 + x^2 - 14x - 6$ divided by $3x + 2$

c) $10x^3 + 11 - 9x^2 - 8x$ divided by $5x - 2$

d) $11x - 4x^4 - 7$ divided by $x - 3$

e) $6x^3 + x^2 + 7x + 3$ divided by $3x + 2$

f) $8x^3 + 4x^2 - 31$ divided by $2x - 3$

g) $6x^2 - 6 + 8x^3$ divided by $4x - 3$

5) The volume, in cubic cm, of a rectangular box can be modelled by the polynomial expression $2x^3 + 17x^2 + 38x + 15$. Determine possible dimensions of the box if the height, in cm, is given by $x + 5$.

6) Determine the value of k such that when $P(x) = kx^3 + 5x^2 - 2x + 3$ is divided by $x + 1$, the remainder is 7.

Answer Key

1) a) 16 b) 31 c) 36

2) a) $x^2 + 2x - 4 + \frac{9}{x+1}$ b) $x^3 + 3x^2 - 2x + 5 = (x+1)(x^2 + 2x - 4) + 9$

3) $x^3 - 2x + 3 + \frac{4}{3x-4}$; check: $3x^4 - 4x^3 - 6x^2 + 17x - 8 = (3x-4)(x^3 - 2x + 3) + 4$

4) a) $x^2 + 5x - 13 + \frac{30}{x+2}$ b) $2x^2 - x - 4 + \frac{2}{3x+2}$ c) $2x^2 - x - 2 + \frac{7}{5x-2}$ d) $-4x^3 - 12x^2 - 36x - 97 - \frac{298}{x-3}$ e) $2x^2 - x + 3 - \frac{3}{3x+2}$ f) $4x^2 + 8x + 12 + \frac{5}{2x-3}$ g) $2x^2 + 3x + \frac{9}{4} + \frac{3}{4(4x-3)}$

5) dimensions: $x + 5$ (height), $x + 3$ (length), $2x + 1$ (width)

6) $k = 3$