

3.4 Solving Exponential Equations Lesson

This document was composed and formatted by Annie Bernatchez.

The course material originated from www.jensenmath.ca

Part 1: Changing the Base of Powers

Exponential functions can be written in many different ways. It is often useful to express an exponential expression using a different base than the one that is given.

Example 1: Express each of the following in terms of a power with a base of 2.

a) 8

b) 4^3

c) $\sqrt{16} \times (\sqrt[5]{32})^3$

d) 12

Rule: $b^{\log_b x} = x$

Part d) shows that any positive number can be expressed as a power of any other positive number.

Example 2: Solve each equation by getting a common base

Remember: if $x^a = x^b$, then $a = b$

a) $4^{x+5} = 64^x$

b) $4^{2x} = 8^{x-3}$

Part 2: Solving Exponential Equations

When you have powers in your equation with different bases and it is difficult to write with the same base, it may be easier to solve by taking the _____ of both sides and applying the _____ of logarithms to remove the variable from the _____.

Example 3: Solve each equation

a) $4^{2x-1} = 3^{x+2}$

Note: Take log of both sides.

Note: Use power law of logarithms.

Note: Use distributive property to expand.

Note: Move variable terms to one side.

Note: Common factor.

Note: Use log rules to isolate the variable and write as a single logarithm.

b) $2^{x+1} = 3^{x-1}$

Part 3: Applying the Quadratic Formula

Sometimes there is no obvious method of solving an exponential equation. If you notice two powers with the same base and an exponent of x , there may be a hidden quadratic.

Example 4: Solve the following equation

$$2^x - 2^{-x} = 4$$

Note: Multiply both sides by 2^x .

Note: Distribute.

Note: Rearrange into standard form $ax^2 + bx + c = 0$.

Let $k = 2^x$ to see the quadratic

Note: Solve using the quadratic formula.

Note: Don't forget to simplify the radical expression.

Now substitute 2^x back in for k and solve

Case 1

Case 2

Part 4: Application Question

Remember:

Equation: $y = a(b)^x$

- a = initial amount
- b = growth ($b > 1$) or decay ($0 < b < 1$) factor
- y = future amount
- x = number of times a has increased or decreased

To calculate x , use the equation:

$$x = \frac{\text{total time}}{\text{time it takes for one growth or decay period}}$$

Example 5: A bacteria culture doubles every 15 minutes. How long will it take for a culture of 20 bacteria to grow to a population of 163 840?

Example 6: One minute after a 100-mg sample of Polonium-218 is placed into a nuclear chamber, only 80-mg remains. What is the half-life of polonium-218?