

3.9 Unit 3 Pre-Test Review – Exponential and Logarithmic Functions

This document was composed and formatted by Annie Bernatchez.

The course material originated from www.jensenmath.ca

1) Sketch a graph of each function. Then, sketch a graph of the inverse of each function. Label each graph with its equation. Also, complete the table of information for each function

a) $f(x) = 2^x$




Table 1. $f(x) = 2^x$

$f(x) =$	$f^{-1}(x) =$
x -int: _____	x -int: _____
y -int: _____	y -int: _____
Domain: _____	Domain: _____
Range: _____	Range: _____
Asymptote: _____	Asymptote: _____

b) $g(x) = \left(\frac{1}{4}\right)^x$





Table 2. $g(x) = \left(\frac{1}{4}\right)^x$

$g(x) =$	$g^{-1}(x) =$
x -int: _____	x -int: _____
y -int: _____	y -int: _____
Domain: _____	Domain: _____
Range: _____	Range: _____
Asymptote: _____	Asymptote: _____

2) State the domain and range for the function, shown below.




Domain: _____


Range: _____

3) Match each graph in the table with the graph of its inverse (A, B, or C). Then write an equation for each function


A)



B)



C)



Graph:		
Equation:		
_____	_____	_____
Letter of Graph of Inverse:		
_____	_____	_____
Equation of inverse:		
_____	_____	_____

4) Rewrite each equation in logarithmic form.

a) $4^3 = 64$

b) $28 = 3^x$

c) $6^3 = y$

d) $512 = 2^9$

5) Rewrite each equation in exponential form.

a) $7 = \log_2 128$

b) $x = \log_b n$

c) $5 = \log_3 243$

d) $19 = \log_b 4$

6) Evaluate without a calculator. Show your work.

Use either:

Rule: if $x^a = x^b$, then $a = b$

Rule: $\log_a(a^b) = b$

a) $\log_2 16$

b) $\log_3 81$

c) $\log_4 \left(\frac{1}{16} \right)$

d) $\log 0.000\,001$

Section 2: 7.4 – Power Law of Logarithms

7) Evaluate each of the following without a calculator using the power law of logarithms.

a) $\log_2 32^3$

b) $\log 1000^{-2}$

c) $\log 0.001^{-1}$

d) $\log_{\frac{1}{4}} \left(\frac{1}{16} \right)^4$

8) Solve for x , correct to 3 decimal places.

a) $x = \log_3 17$

b) $\log_2 0.35 = x$

c) $4^x = 10$

d) $80 = 100\left(\frac{1}{2}\right)^x$

Section 3: 7.3 – Product and Quotient Laws of Logarithms

9) Use the change of base formula to evaluate. Round to one decimal place.

a) $\log_9 12$

b) $\log_{0.25} 52$

10) Write as a single logarithm. Then evaluate without a calculator.

a) $\frac{\log 16}{\log 4}$

b) $\frac{\log\left(\frac{8}{27}\right)}{\log\left(\frac{2}{3}\right)}$

11) Solve, to two decimal places

a) $\log 4^x = 7$

b) $12 = \log_3 4^m$

12) An investment earns 12% interest, compounded annually. The amount, A , that the investment is worth as a function of time, t , in years, is given by $A = 1500(1.12)^t$. Use the equation to determine...

a) the value of the investment after 4 years

b) how long it will take for the investment to double in value

13) Write as a single logarithm

a) $\log_7 8 + \log_7 4 - \log_7 16$

b) $2\log a + \log(3b) - \frac{1}{2}\log c$

14) Write as a sum or difference of logarithms. Simplify if possible.

a) $\log(a^2bc)$

b) $\log\left(\frac{k}{\sqrt{m}}\right)$

15) Evaluate, using the laws of logarithms.

a) $\log_6 8 + \log_6 27$

b) $\log_4 128 - \log_4 8$

c) $2\log 2 + 2\log 5$

d) $2\log 3 + \log\left(\frac{25}{2}\right)$

16) Simplify

a) $\log(2m + 6) - \log(m^2 - 9)$

b) $\log(x^2 + 2x - 15) - \log(x^2 - 7x + 12)$

Section 4: 7.1/7.2 – Solving Exponential Equations

17) Write each as a power of 4

a) 64

b) $\frac{1}{16}$

c) $(\sqrt[3]{8})^5$

18) Write 20 as a power of 5.

19) Solve each equation

a) $3^{5x} = 27^{x-1}$

b) $8^{2x+1} = 32^{x-1}$

20) Solve exactly. Then use your calculator to evaluate correct to 3 decimal places.

a) $3^{x-2} = 5^x$

b) $2^{k-2} = 3^{k+1}$

21) Solve the following equations; round to 2 decimal places where appropriate.

a) $3^x = 12$

b) $10 = 2 \cdot 4^{x+2}$

c) $3^x = 4^{1-x}$

22) Solve each equation. Check for extraneous roots.

a) $4^{2x} - 4^x - 20 = 0$

Let $k = 4^x$

Case 1

Case 2

b) $2^x + 12(2)^{-x} = 7$

Let $k = 2^x$

Case 1

Case 2

Section 5: 7.4 – Solving Logarithmic Equations

23) Solve each equation

a) $\log_4 x = 1.8$

b) $\log_5 x - \log_5(x - 2) = 1$

c) $5^{2x} = 2(5)^x + 1$

Let $k = 5^x$

Case 1

Case 2

24) Solve

a) $\log(2x + 10) = 2$

b) $1 - \log(2x) = 0$

25) Solve. Check for extraneous roots.

a) $\log_2 x + \log_2(x + 2) = 3$

b) $\log_3(3x + 7) = 2$

c) $\log_5(2x + 1) = 1 - \log_5(x + 2)$

Section 6: 7.4 – Applications

Table 3. *Exponential Formulas*

General	Half-life	Doubling
$A(t) = A_0(1 + i)^t$	$A(t) = A_0\left(\frac{1}{2}\right)^{\frac{t}{H}}$	$A(t) = A_0(2)^{\frac{t}{D}}$
i is percent growth (+) or decay (-)	H is the half-life period	D is the doubling period

Table 4. *Logarithmic Formulas*

pH	Decibel	Richter
$pH = -\log[H^+]$	$\beta_2 - \beta_1 = 10\log\left(\frac{I_2}{I_1}\right)$	$M = \log\left(\frac{I}{I_0}\right)$
pH is acidity and $[H^+]$ is concentration of hydronium ions in mol/L	β is loudness in dB and I is intensity of sound in W/m^2	M is magnitude measured by richters, I is intensity

26) When you drink a cup of coffee or a glass of cola, or when you eat a chocolate bar, the percent, P , of caffeine remaining in your bloodstream is related to the elapsed time, t , in hours by $t = 5\left(\frac{\log P}{\log 0.5}\right)$

a) How long will it take for the amount of caffeine to drop to 20% of the amount consumed?

b) Suppose you drink a cup of coffee at 9:00 am, what percent of the caffeine will remain in your body at noon?

27) A 50-mg sample of cobalt-60 decays to 40 mg after 1.6 minutes.

a) Determine the half-life of cobalt-60.

b) How long will it take for the sample to decay to 5% of its initial amount?

28) Determine the pH, correct to one decimal place, of a solution with each hydronium ion concentration.

a) 0.000 316 mol/L

b) 7.9×10^{-9} mol/L

29) Calculate the hydronium ion concentration, correct to two decimal places, if the pH of a solution is

a) 2.2

b) 11.6

30) Use the sound level scale in your notes to answer the following:

a) How many times as intense is a normal conversation compared to a whisper?

b) How many times as intense is normal city traffic compared to a shout?

31) The intensity of sound in a library is estimated to be one thousandth that of normal conversation. What is the decibel rating for the library?

32) How many times as intense is an earthquake with a magnitude of 7.2 than an earthquake with a magnitude of 5.6?

33) If an earthquake is 390 times as intense as an earthquake with a magnitude of 4.2 on the Richter scale, what is the magnitude of the more intense earthquake?

34) The absolute magnitude of star A is -4.5 and that of star B is 0.2. How many times as bright is star A than star B, to the nearest unit?

35) An altimeter is a device that measures the height of a plane above the ground. It works based on air pressure according to the formula $h = 18\,400 \log \frac{P_0}{P}$, where h is the height above the ground in metres, P is the air pressure at that height, and P_0 was the air pressure on the ground at takeoff. Air pressure is measured in kilopascals (kPa).

a) Air pressure on the ground was 102 kPa. If the airplane instruments measure a pressure of 32.5 kPa outside

the plane, what is the height of the airplane to the nearest metre?

b) What is the outside air pressure for a plane flying at 11 000 metres? Assume a ground pressure of 102.5 kPa. Round to one decimal place.

c) How high would a plane have to be flying when it encountered air pressure in the air that was half the air pressure on the ground? Round to the nearest metre.