
5.5 Cross Product of Vectors Lesson

This document was composed and formatted by Annie Bernatchez.


The course material originated from www.jensenmath.ca

The cross product of two vectors $\vec{a}$ and $\vec{b}$ in $\mathbb{R}^3$ is the vector that is **perpendicular** to these vectors such that the vectors $\vec{a}$, $\vec{b}$, and $\vec{a}\times\vec{b}$ form a right-handed system.

Right handed system tells you to point your hand along vector $\vec{a}$ and curl your fingers towards vector $\vec{b}$. Your thumb will be pointing in the direction of $\vec{a}\times\vec{b}$. Notice that $\vec{a}\times\vec{b}$ and $\vec{b}\times\vec{a}$ point in opposite directions.



Sometimes the direction of the cross product is defined by either 'in to the page' or 'out of the page':




$\vec{a}\times\vec{b}$	$\vec{b}\times\vec{a}$

3D visualization:


www.geogebra.org/3d/jyqcr3bf

Copyright © 2026 Annie Bernatchez—All rights reserved.



Properties of Cross Product:

- $\vec{u}\times\vec{v} = -(\vec{v}\times\vec{u})$
- $\vec{u}\times(\vec{v}+\vec{w}) = \vec{u}\times\vec{v} + \vec{u}\times\vec{w}$
- $(\vec{u}+\vec{v})\times\vec{w} = \vec{u}\times\vec{w} + \vec{v}\times\vec{w}$
- If $\vec{u}\times\vec{v} = 0$, and $\vec{u}$ and $\vec{v}$ are non-zero, then $\vec{u}$ and $\vec{v}$ are collinear.
- $k(\vec{u}\times\vec{v}) = (k\vec{u})\times\vec{v} = \vec{u}\times(k\vec{v})$
- $|\vec{u}\times\vec{v}| =$ the area of the parallelogram defined by $\vec{u}$ and $\vec{v}$



Part 2: Cross Product of Geometric Vectors


Formula: $\vec{a}\times\vec{b} = (|\vec{a}||\vec{b}|\sin\theta)\hat{n}$

θ is the angle between the vectors

$\hat{n}$ is a unit vector perpendicular to both $\vec{a}$ and $\vec{b}$

Copyright © 2026 Annie Bernatchez—All rights reserved.

Example 1: If $|\vec{u}| = 30$, $|\vec{v}| = 20$, the angle between $\vec{u}$ and $\vec{v}$ is 40°, and $\vec{u}$ and $\vec{v}$ are in the plane of the page, find…

a) $\vec{u} \times \vec{v}$ b) $\vec{v} \times \vec{u}$	

Part 3: Cross Product of Algebraic Vectors


Formula:

If $\vec{a} = [a_1, a_2, a_3]$ and $\vec{b} = [b_1, b_2, b_3]$

$\vec{a} \times \vec{b} = [a_2b_3 - a_3b_2,\ a_3b_1 - a_1b_3,\ a_1b_2 - a_2b_1]$

$= (a_2b_3 - a_3b_2)\vec{i} + (a_3b_1 - a_1b_3)\vec{j} + (a_1b_2 - a_2b_1)\vec{k}$

How to set it up:



Example 2: If $\vec{p} = [-1, 3, 2]$ and $\vec{q} = [2, -5, 6]$, calculate each of the following:


a) $\vec{p} \times \vec{q}$

Copyright © 2026 Annie Bernatchez—All rights reserved.

b) $\vec{q} \times \vec{p}$

Example 3a: Determine the area of the parallelogram defined by the vectors $\vec{u} = [4, 5, 2]$ and $\vec{v} = [3, 2, 7]$.

Example 3b: Determine the angle between the vectors $\vec{u}$ and $\vec{v}$.



Copyright © 2026 Annie Bernatchez—All rights reserved.