

6.7 Intersection of Planes Lesson with solutions

This document was composed and formatted by Annie Bernatchez.

The course material originated from www.jensenmath.ca

Part 1: Intersection of 2 Planes

Line of Intersection	Infinite Solutions	No Solutions
		
If 2 distinct planes intersect, the solution is the set of points that lie on the line of intersection.	If the planes are coincident, every point on the plane is a solution	Parallel and distinct planes do not intersect

When solving a system of 2 planes, check if the planes are parallel by analyzing their normal vectors. If they are parallel, determine if they are coincident (infinite solutions) or distinct (no solutions). If the normals are not parallel, that means the planes are not parallel either. You can find the line of intersection by:

i) Eliminating a variable using the method of elimination

ii) Choose 1 of the remaining variables to be the parameter t

iii) Write the other 2 variables in terms of the parameter t as well to get the parametric equation of the line of intersection

Example 1: Describe how the planes in each pair intersect.

a) $\pi_1 : 2x - y + z - 1 = 0$

$\pi_2 : x + y + z - 6 = 0$

$\vec{n_1} = [2, \ -1, \ 1]$, $\vec{n_2} = [1, \ 1, \ 1]$

$\vec{n_1} \neq k\vec{n_2}$

The normals are not parallel and therefore the planes are not parallel. This means they intersect in a line.

① $\quad 2x - y + z - 1 = 0$

② $\quad x + y + z - 6 = 0$

① + ② : $\quad 3x + 2z - 7 = 0$

Assign a variable to be the parameter t. Let $z = t$ and then write the other two variables in terms of t.

$3x + 2t - 7 = 0$

$x = -\frac{2}{3}t + \frac{7}{3}$

Copyright © 2026 Annie Bernatchez—All rights reserved.

$\left(-\frac{2}{3}t+\frac{7}{3}\right)+y+t-6=0$

$\frac{1}{3}t+y-\frac{11}{3}=0$

$y=-\frac{1}{3}t+\frac{11}{3}$

The parametric equations of the line of intersection are: $\ell:\begin{cases}x=-\frac{2}{3}t+\frac{7}{3}\\ y=-\frac{1}{3}t+\frac{11}{3}\\ z=t\end{cases}$

Note: the direction vector was simplified

Vector equation is: $[x,\ y,\ z]=\left[\frac{7}{3},\ \frac{11}{3},\ 0\right]+t[-2,\ -1,\ 3]$

b) $\pi_3: 2x-6y+4z-7=0$

$\pi_4: 3x-9y+6z-2=0$

$\vec{n_3}=[2,\ -6,\ 4]$, $\vec{n_4}=[3,\ -9,\ 6]$

$1.5\vec{n_3}=\vec{n_4}$

The normals are parallel. Therefore, the planes are parallel. They are either coincident or distinct. Solve using elimination to see if you get infinitely many solutions or no solutions.

$3\times③:\quad 6x-18y+12z=21$

$2\times④:\quad 6x-18y+12z=4$

$0=17$

This equation is never true. Therefore, there are no solutions and the planes are parallel and distinct.

c) $\pi_5: x+y-2z+2=0$

$\pi_6: 2x+2y-4z+4=0$

$\vec{n_5}=[1,\ 1,\ -2]$, $\vec{n_6}=[2,\ 2,\ -4]$

$2\vec{n_5}=\vec{n_6}$

The normals are parallel. Therefore, the planes are parallel. They are either coincident or distinct. Solve using elimination to see if you get infinitely many solutions or no solutions.

$2\times⑤:\quad 2x+2y-4z=-4$

$⑥:\quad 2x+2y-4z=-4$

$0=0$



This equation is always true. Therefore, there are infinitely many solutions. The planes are coincident.

Part 2: Intersection of 3 Planes

A system of three planes is consistent if it has one or more solutions. A system of three planes is inconsistent if it has no solution.

3 scenarios of consistent solutions:

Copyright © 2026 Annie Bernatchez—All rights reserved.

Intersect at 1 Point	Intersect in a Line	Infinite Solutions
		
Normals are not parallel and not coplanar.	Normals are not parallel but they are coplanar.	Normals are parallel and equations are scalar multiples of eachother.

Normals are parallel if each one is a scalar multiple of the others.

Normals are coplanar if $\vec{n_1} \cdot \vec{n_2} \times \vec{n_3} = 0$

Example 2: For each set of planes, describe the number of solutions and how the planes intersect.

a) $\pi_1 : 2x + y + 6z - 7 = 0$

$\pi_2 : 3x + 4y + 3z + 8 = 0$

$\pi_3 : x - 2y - 4z - 9 = 0$

Normals are not parallel. Therefore, they intersect at a point or in a line.

Eliminate a variable using two different pairs of equations:

① $2x + y + 6z = 7$

$2\times$② $6x + 8y + 6z = -16$

$-4x - 7y = 23$

$2\times$① $4x + 2y + 12z = 14$

$3\times$③ $3x - 6y - 12z = 27$

$7x - 4y = 41$

$7\times(① - ②):$ $-28x - 49y = 161$

$4\times(① + ③):$ $28x - 16y = 164$

$-65y = 325$

$y = -5$

Now that you have a unique solution for one variable, you can sub back in to previous equations to solve for the remaining variables:

$-4x - 7(-5) = 23$

$x = 3$

$2x + y + 6z = 7:$ $2(3) + (-5) + 6z = 7$

$z = 1$

The three planes form a consistent system. The point of intersection is $(3, -5, 1)$.

b) $\pi_4 : x - 5y + 2z - 10 = 0$

$\pi_5 : x + 7y - 2z + 6 = 0$

Copyright © 2026 Annie Bernatchez—All rights reserved.

$\pi_6 : 8x + 5y + z - 20 = 0$

Normals are not parallel. Therefore, they intersect at a point or in a line.

Eliminate a variable using two different pairs of equations:

④ $x - 5y + 2z = 10$

⑤ $x + 7y - 2z = -6$

$2x + 2y = 4$

$2\times$⑥ $16x + 10y + 2z = 40$

$-15x - 15y = -30$

$-\frac{1}{15}\times(④ - ⑥):$ $x + y = 2$

$\frac{1}{2}\times(④ + ⑤):$ $x + y = 2$

$0 = 0$

This means that there are infinitely many values of x and y that satisfy the equations. Therefore, they intersect in a line. Write the parametric equations of the line. Start by letting a variable equal a parameter:

$x = t$

$x + y = 2:$ $t + y = 2$

$y = 2 - t$

$x - 5y + 2z = 10:$ $t - 5(2 - t) + 2z = 10$

$t - 10 + 5t + 2z = 10$




$2z = -6t + 20$

$z = -3t + 10$

Parametric Equations: $\ell : \begin{cases} x = t \\ y = 2 - t \\ z = 10 - 3t \end{cases}$

Vector Equation: $[x, y, z] = [0, 2, 10] + t[1, -1, -3]$

3 scenarios of inconsistent solutions:

2 planes are parallel and the third intersects both of the parallel planes	**The planes intersect in pairs**	**The planes are parallel and at least two are distinct**
		
Two normals are parallel but the third is not	Normals are not parallel but they are coplanar.	Normals are parallel but the equations are not scalar multiples of each other

Example 3: For each set of planes, describe the number of solutions and how the planes intersect.

Copyright © 2026 Annie Bernatchez—All rights reserved.

Example 3: For each set of planes, describe the number of solutions and how the planes intersect.

a) $\pi_1 : 3x + y - 2z = 12$

$\pi_2 : 3x - 5y + z = 8$

$\pi_3 : 12x + 4y - 8z = -4$

π_1 and π_3 are parallel since $4\vec{n_1} = \vec{n_3}$ but π_2 is not parallel to them. $\pi_1 \neq k\pi_3$, so they are parallel but distinct. This is an inconsistent system with two parallel but distinct planes that are intersected by a third plane.

b) $\pi_4 : x + 3y - z = -10$

$\pi_5 : 2x + y + z = 8$

$\pi_6 : x - 2y + 2z = -4$

None of the normals are scalar multiples of each other so none of the planes are parallel. They either intersect at a point, in a line, or not at all. Solve using elimination to determine which scenario it is:

④ $x + 3y - z = -10$

⑤ $2x + y + z = 8$

$3x + 4y = -2$

$2\times$④ $2x + 6y - 2z = -20$

⑥ $x - 2y + 2z = -4$

$3x + 4y = -24$

(④ + ⑤) : $3x + 4y = -2$

(④ + ⑥) : $3x + 4y = -24$

$0 = 22$

There are no solutions to this equation. Therefore, this is an inconsistent system with no solutions. The planes must intersect in pairs.

c) $\pi_7 : 4x - 2y + 6z = 35$

$\pi_8 : -10x + 5y - 15z = 20$

$\pi_9 : 6x - 3y + 9z = -50$

Notice that the normals between each pair of planes are constant multiples of each other. Therefore, the planes are all parallel. However, none of the planes are scalar multiples of each other so they are all parallel but distinct. There are no points of intersection.

Copyright © 2026 Annie Bernatchez—All rights reserved.